



North Dakota Public Health Association 2022 Resolution to Use Strategies to Reduce Alcohol-related Harm

Whereas North Dakota is the leading binge drinking state measured by the largest number of drinks consumed on an occasion in the past 30 days among adults who reported binge drinking (5.9), and is still the 3rd leading binge drinking state among adults, with a prevalence of 22.7% of adults aged 18 or greater, trailing only Iowa and Wisconsin¹;

Whereas North Dakota is a state that values life, and yet North Dakota ranks as the #47 worst state for alcohol deaths, with 17.2 deaths per 100,000, which is twice the national average of 9.9; and this number has increased by 58% from 2005 (10.9) to 2018 (17.2) ¹;

Whereas alcohol is responsible for substantial societal and personal harm,^{2,3} and chronic excessive alcohol use contributes directly to cardiovascular disease,⁴ nutritional deficiency,⁵ cancer,⁶ accelerated aging,⁷ gastrointestinal deaths, liver cirrhosis in younger adults,⁸ injuries, and alcohol use disorders²;

Whereas among all U.S. cirrhosis deaths in 2015, approximately 49.5% were related to alcohol use and occurred in younger adults ages 25-34,⁹ so this is a health concern for all ages;

Whereas there is increasing evidence that there is no level of alcohol intake that is considered beneficial to health¹⁰;

Whereas underage drinking is a common problem, including businesses failing compliance checks regarding selling alcohol to minors¹¹⁻¹³;

Whereas alcohol remains low cost and is become more affordable, not less,¹⁴ despite global evidence that alcohol per capita consumption is associated with affordability,¹⁵ and the alcohol industry indiscriminately promotes its products to increase sales, without being held responsible for the personal and societal harm caused by alcohol misuse¹⁶;

Whereas persistently high levels of alcohol harm create a heavy social, economic, and health burden. For example, one in every 10 deaths in the European region is caused by alcohol every year, and nearly one in every four deaths among young adults aged 20–24 years is due to alcohol products¹⁷;

Whereas the COVID-19 pandemic resulted in loss of purpose for many people, loss of jobs, and being required to work from home, so that drinking during the day increased; and it became easier to purchase alcohol, including curbside pickup, resulting in a 25% increase in alcohol-related deaths in 2020¹⁸;

Whereas alcohol is a psychoactive substance and a known carcinogen, which causes substantial harms to alcohol users, their families, communities, as well as societies and economies in general, it is far less regulated than tobacco, including the regulations of the level of taxation;

Whereas alcohol consumption is considered a social norm, resulting in many people in society resisting any restrictions on alcohol, so that alcohol taxes have been under-utilized as a public health measure. North Dakota's excise tax is \$0.50/gallon for wine, \$0.16/gallon for beer, and \$2.50/gallon for liquor, ranking 32nd, 34th, and 28th in the United States¹⁹;

Whereas the state of Washington has the highest alcohol excise tax, with \$33.22 per gallon for liquor; and is among the states with the lowest prevalence of binge drinking (15.4%), and the lowest number of drinks consumed when drinking (5.3) ¹⁹;

Whereas the CDC Community Preventive Services Task Force has ranked increasing alcohol excise taxes to prevent excessive alcohol use and other harms as an evidence-based best practice, with the ability to decrease total alcohol consumption by 7.7% for every 10% increase in price²⁰;

Whereas people consistently value health as the most important condition for a happy and flourishing life,²¹ and while alcohol is known to increase the feeling of social cohesiveness; in fact, excess alcohol consumption compromises health, happiness and human flourishing to a great degree; and

Whereas Governor Doug and First Lady Kathryn Burgum have established the Recovery Reinvented program to eliminate the shame and stigma of addiction in North Dakota and to help all persons in North Dakota to prevent chemical misuse from beginning, treat addiction for all persons seeking help, and support a life of recovery; now, therefore, be it

Resolved, that the North Dakota Public Health Association, on behalf of its members recommends:

1. increase the excise tax on beer in bottles and cans from \$0.16/gallon to \$0.26/gallon, wine from \$0.50/gallon to \$0.87/gallon, and liquor from \$2.50/gallon to \$14.27/gallon; in addition to all federal excise taxes;
2. the North Dakota legislature conduct a study to determine a cost recovery approach to cover the costs of alcohol harms to society;
3. the legislature to make alcohol-related policy decisions from a Health in All Policies perspective.
4. decrease the density of alcohol sales outlets;
5. reduce the hours and the days of alcohol sales;
6. forbid direct to consumer advertising of alcohol products;
7. increase the visibility of "alcohol can cause serious harm" messaging on alcohol product packaging;
8. require Responsible Beverage Service (RBS) training as a condition of on-sale alcohol sale licensure; and
9. fund alcohol compliance checks to reduce youth access to alcohol.

References:

1. Centers for Disease Control and Prevention. Data on Excessive Drinking. Centers for Disease Control and Prevention. <https://www.cdc.gov/alcohol/data-stats.htm>
2. Rehm J, Mathers C, Popova S, Thavorncharoensap M, Teerawattananon Y, Patra J. Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *Lancet*. 2009;373:2223-33.
3. Rehm J, Room R, Graham K, Monteiro M, Gmel G, Sempos C. The relationship of average volume of alcohol consumption and patterns of drinking to burden of disease. *Addiction*. 2003;98:1209–28.
4. Zaridze D, Brennan P, Boreham J, et al. Alcohol and cause-specific mortality in Russia: a retrospective case-control study of 48 557 adult deaths. *Lancet*. 2009;373:2201–14.
5. Zahr NM, Sullivan EV, Rohlfing T, et al. Concomitants of alcoholism: differential effects of thiamine deficiency, liver damage, and food deprivation on the rat brain in vivo. *Psychopharmacology*. 2016;233(14):2675-2686.
6. Rehm J, Shield KD, Weiderpass E. Alcohol consumption. A leading risk factor for cancer. *Chemico-Biological Interactions*. 2020;331:109280.
7. Sullivan EV, Pfefferbaum A. Brain-behavior relations and effects of aging and common comorbidities in alcohol use disorder: A review. *Neuropsychology*. 2019;33(6):760.
8. Yoon Y-H, Chen CM. Surveillance Report# 111: Liver Cirrhosis Mortality in the United States: National, State, and Regional Trends, 2000–2015. *Rockville, MD: National Institute on Alcohol Abuse and Alcoholism*. 2018;
9. Chung W, Jo C, Chung WJ, Kim DJ. Liver cirrhosis and cancer: comparison of mortality. *Hepatology international*. 2018;12(3):269-276.
10. Burton R, Sheron N. No level of alcohol consumption improves health. *The Lancet*. 2018;392(10152):987-988.
11. Toomey TL, Komro KA, Oakes JM, Lenk KM. Propensity for illegal alcohol sales to underage youth in Chicago. *Journal of community health*. 2008;33(3):134-138.
12. van Hoof JJ, Gosselt JF. Underage alcohol sales—It only takes a minute: A new approach to underage alcohol availability. *Journal of studies on alcohol and drugs*. 2013;74(3):423-427.
13. Freisthler B, Gruenewald PJ, Treno AJ, Lee J. Evaluating alcohol access and the alcohol environment in neighborhood areas. *Alcoholism: clinical and experimental research*. 2003;27(3):477-484.
14. Knai C, Petticrew M, Durand MA, Eastmure E, Mays N. Are the Public Health Responsibility Deal alcohol pledges likely to improve public health? An evidence synthesis. *Addiction*. 2015;110(8):1232-1246.
15. Casswell S, Huckle T, Parker K, Romeo J, Graydon-Guy T. Benchmarking alcohol policy based on stringency and impact: The International Alcohol Control (IAC) policy index. *PLOS Global Public Health*. 2022;2(4):e0000109. doi:PLOS Global Public Health 2(4): e0000109. <https://doi.org/10.1371/journal.pgph.0000109>
16. Jernigan D, Ross CS. The Alcohol Marketing Landscape: Alcohol Industry Size, Structure, Strategies, and Public Health Responses. *Journal of Studies on Alcohol and Drugs, Supplement*. 2020/03/01 2020;(s19):13-25. doi:10.15288/jsads.2020.s19.13
17. Neufeld M, Rovira P, Ferreira-Borges C, et al. Impact of introducing a minimum alcohol tax share in retail prices on alcohol-attributable mortality in the WHO European Region: A modelling study. *The Lancet Regional Health – Europe*. 2022;p.100325.
18. White AM, Castle IJP, Powell PA, Hingson RW, Koob GF. Alcohol-Related Deaths During the COVID-19 Pandemic. *JAMA*. 2022;doi:10.1001/jama.2022.4308
19. World Population Review. Alcohol Tax by State 2022. April 22, 2022, <https://worldpopulationreview.com/state-rankings/alcohol-tax-by-state>
20. The Community Preventive Services Task Force. *Alcohol -- Excessive Consumption: Increasing Alcohol Taxes*. 2007. Accessed May 3, 2022. <https://www.thecommunityguide.org/findings/alcohol-excessive-consumption-increasing-alcohol-taxes>
21. Levin J. Human flourishing: a new concept for preventive medicine. *American Journal of Preventive Medicine*. 2021;61(5):761-764.